

# The biology and therapeutic promise of small RNAs

Phillip Sharp

*Koch Institute for  
Integrative Cancer Research*

# RNA in the transfer of information from DNA to protein

---


- a) mRNA - message from DNA
- b) t-RNA - intermediate between nucleic acid sequence and amino acid
- c) Ribosomal RNA - components of decoding machine

Cellular environment can  
modify nature of mRNA in transfer  
of information from DNA to protein

---

- a) Alternative RNA splicing
- b) Alternative polyadenylation  
and RNA editing

EST evidence has demonstrated several modes of variation for transcripts coming from a single locus




- Standard transcriptional activation
- Alternative promoter usage
- Exon inclusion/exclusion
- 3' UTR utilization
- All of the above


# *RNA interference (RNAi)*

- Generated by dsRNA
- Silences genes at post-transcriptional stage by cleavage of mRNA
- Related process silences genes at translation
- Related process silences genes at transcriptional stage: transposable elements in genome

# Short RNAs silence by mRNA degradation



# Liposomal formulations for systemic RNAi




- Liposomes are organized lipid nanoparticles used in pharmaceutical drug delivery
  - Used in over 12 approved drugs
- Enables delivery of siRNAs into cells
  - Highly efficient for liver delivery

# *In vivo* silencing of 2 distinct genes


Mouse

Delivery of 2 siRNAs in liposomal nanoparticles

**FVII  
Liver  
mRNA**



**ApoB  
Liver  
mRNA**







# Repeat silencing of Factor VII

## Rat model

Repeat dosing over 3 months is highly effective



# Prolonged survival with ALN-VSP treatment



Orthotopic Tumor Models (Hep3B)

1. Treated 18 days post seeding; IV bolus injections of 4 mg/kg VSP or control siRNA 2x/wk for 3 wks
2. Treated 26 days post seeding; IV bolus injections of 4 mg/kg VSP or control siRNA 2x/wk for 3 wks

The discovery of 250 to 1000 new genes that encode microRNAs. These probably regulate 25-50% of all genes in vertebrates.

V. Ambros (Lee *et al.* 1993)


L. Ruvkun (Wightman *et al.* 1993)

T. Tuschl (Lagos-Quintana *et al.* 2001)

D. Bartel (Lau *et al.* 2001)


V. Ambros (Lee and Ambros, 2001)

# microRNA pathway



# Proposed anatomy of miRNA/mRNA interactions


---




V. Ambros (Lee *et al.* 1993)

L. Ruvkun (Wightman *et al.* 1993)

# Structural insights into mammalian Argonaute function from studies of bacterial Argonaute (D. Patel and T. Tuschl)




# *T. th* Ago-Guide DNA-Target RNA Ternary Complex





# Conserved seed matches in 3' UTRs indicate extensive regulation by miRNAs

---



25-50% of all mRNAs  
interact with miRNAs

B.P. Lewis *et al.* 2005  
Cell 120:15



# Summary of target predictions for miRNA regulation in mammals


---

87 evolutionarily conserved “seed families” of miRNAs



Distribution of preferentially conserved target sites:

- 45% of genes - no sites in 3' UTR
- 55% of genes - one or more target sites
  - Average is 4.2 sites per UTR and more than one family
  - Average 500 target mRNAs per family

# microRNA regulation in cancer network



# let-7 is negatively regulated in non-small cell lung cancer (NSCLC)



|                 | Months after Surgery |    |    |    |
|-----------------|----------------------|----|----|----|
| Numbers at risk |                      |    |    |    |
| Cluster 1       | 67                   | 53 | 41 | 21 |
| Cluster 2       | 76                   | 71 | 63 | 45 |

# A role of microRNA regulation-Robustness

---

## 1) Stabilize quiescent and differentiated state

- microRNA-let7 suppresses Ras, etc (growth) (*Frank Slack*)
- Cell-type specific microRNAs suppresses genes expressed in other cellular states (*Lee Lim-05 and Stephen M. Cohen-05*)
- Suppress cell-type specific genes

## 2) Control responses to changes in environment

- Protection against abnormal cellular states (p53 and NF- $\kappa$ B) and stabilize cellular states from transient “stress”

# Alternative cleavage and polyadenylation

---

## **Proliferating Cells Express mRNAs with Shortened 3' Untranslated Regions and Fewer MicroRNA Target Sites**

Rickard Sandberg,<sup>1\*†</sup> Joel R. Neilson,<sup>2\*</sup> Arup Sarma,<sup>3</sup>  
Phillip A. Sharp,<sup>1,2‡</sup> Christopher B. Burge<sup>1‡</sup>

*Science* 20 June 2008:  
Vol. 320. no. 5883, pp. 1643

## **Widespread Shortening of 3'UTRs by Alternative Cleavage and Polyadenylation Activates Oncogenes in Cancer Cells**

Christine Mayr<sup>1,2,3,4,\*</sup> and David P. Bartel<sup>1,2,3,\*</sup>

*Cell* 138, 673–684, August 21, 2009

## **Progressive lengthening of 3' untranslated regions of mRNAs by alternative polyadenylation during mouse embryonic development**

Zhe Ji<sup>1</sup>, Ju Youn Lee<sup>1</sup>, Zhenhua Pan<sup>1</sup>, Bingjun Jiang, and Bin Tian<sup>2</sup>


7028–7033 | PNAS | April 28, 2009 | vol. 106

Department of Biochemistry and Molecular Biology, New Jersey Medical School, University of Medicine and Dentistry of New Jersey, Newark, NJ 07103

# A model for evasion of post-transcriptional regulation


## Regulators of translation and degradation

- RNA binding proteins
- miRNAs





# Post-transcriptional regulation critical for quiescent tissue




| Quiescent tissue | Proliferative tissue | Tumor tissue | Cultured cell line |
|------------------|----------------------|--------------|--------------------|
|                  |                      |              |                    |
| +++              | +                    | —            | —                  |
| +                | ++                   | ++           | +                  |
| —                | +                    | ++           | +++                |
| ++++             | ++++                 | ++++         | ++++               |

Gene structure and isoforms

Relative expression

# ES cells survive Dicer deletion, but are no longer pluripotent

---



Genes Dev. 2005 Feb 15;19(4):489-501

“However, DCR<sup>-/-</sup> cells did not contribute to the generation of chimeric mice when injected into blastocysts, unlike DCR<sup>neo/+</sup> and DCR<sup>neo/neo</sup> cells, which gave rise to high-percentage chimeras (data not shown). Furthermore, the Dicer-deficient cells failed to generate detectable teratomas upon subcutaneous injection into nude mice ([Fig. 2C](#)).”



# Generation of Dicer null sarcoma cell line

\* **Kras<sup>G12D</sup>, Trp53<sup>-/-</sup>, Dicer1<sup>flox/-</sup>**  
murine sarcoma cells



250 nM Tamoxifen,  
24 hr




1:5000 clonal  
plating



Day 7  
select colonies





Genotype




\*Generated by Madhu  
Kumar and Tyler Jacks

# Dicer null cells retain the ability to form tumors

## Subcutaneous tumors in nude mice



*Dicer1*<sup>flox/-</sup>



*Dicer1*<sup>-/-</sup>

# Literature on microRNA and Stress


## Genetic Data

Knockout of miR-14 in *Drosophila* results in sensitivity to salt stress and reduced lifespan (Xu et al., Curr Biol 2003)

Heart-specific miR-208 regulates stress-dependent cardiac growth in mice (Rooij et al., Science 2007)

miR-8 regulates the response to osmotic stress in zebrafish embryos (Flynt et al., JCB 2009)

miR-7 stabilizes gene regulatory networks against environmental fluctuations during *Drosophila* development (Li et al., Cell 2009)




# Literature on microRNA and Stress


## Localization

Upon oxidative stress or inhibition of translation initiation, we found that Argonaute localizes to Stress Granules (SGs) in a miRNA-dependent manner.

Argonaute at SGs continues to exchange with the cytoplasm, in which 50% of Argonaute at SGs are replaced within 20 seconds.  
(Leung et al., PNAS 2006)



# Stressed cells form cytoplasmic stress granules (SGs) in which mRNA translation is arrested in pre-initiation complex




## Stress Granules (SGs)

- first identified in plant
- conserved from worm to human
- stalled initiation complex
- enriched with RNA binding proteins (TIA-1, G3BP-1, PABP-1, and Ago2)
- proposed role of mRNA triage (degradation, reinitiation and storage)

modified from  
Hilliker and Parker  
Nat Cell Biol 2008

# Dynamic localization of post-transcriptional regulators

microRNA-binding factor Argonaute (Ago2)



EGFP-Ago2


Time 0: P-bodies with Ago2 -  
degradative sites for  
mRNAs

**Stress granules forms upon**  
inhibition of translation initiation  
heat shock  
UV irradiation  
viral infection  
energy depletion  
proteasomeinhibition  
hypoglycemia  
hypoxia

Leung et al., PNAS  
2006

# Dynamic localization of post-transcriptional regulators

microRNA-binding factor Argonaute (Ago2)



EGFP-Ago2

Time 0: P-bodies with Ago2 -  
degradative sites for  
mRNAs



Time 35: Formation of stress  
granules containing Ago2

**Stress granules forms upon**  
inhibition of translation initiation  
heat shock  
UV irradiation  
viral infection  
energy depletion  
proteasomeinhibition  
hypoglycemia  
hypoxia

Leung et al., PNAS  
2006





# Stress Granules present in Hypoxic tumor cells





# Evidence that Poly(ADP-ribose), PAR, is matrix for SG formation and influences miRNA regulation of translation

## Role of PAR in DNA damage



### Poly-ADP-ribose (PAR) modified

- upon DNA damage
- during mitosis at mitotic spindle


modified from Schreiber et al  
Nat Rev Mol Cell Biol 2006

Poly(ADP-ribose) is **essential** in multicellular eukaryotes


- ✿ Required for Chromosome Integrity, Transcription and Cell Division
- ✿ PARP-1 is targeted by pharmaceutical companies for BRCA1- Cancer
- ✿ Inhibitors have been beneficial in ischemia, inflammation, degenerative and vascular diseases

# Proposed model of Stress Granule formation

---




# Proposed model of Stress Granule formation



# Proposed model of gene regulation by Poly(ADP-ribose)

---



# Conclusions

---

- PAR, 5 PARPs and 2 PARG isoforms are localized in RNA-binding protein enriched-Stress Granules.
- Poly(ADP-ribose) is required for structural integrity of Stress Granules.
- Specific RNA-binding proteins are modified by PAR upon stress, effecting post-transcriptional gene regulation.
- Under stress conditions, the level of miRNA regulation is reduced.
- Poly(ADP-ribose) probably regulates translation under normal and stress conditions.

# A role of microRNA regulation-Robustness

---

## 1) Stabilize quiescent and differentiated state

- microRNA-let7 suppresses Ras, etc (growth) (*Frank Slack*)
- Cell-type specific microRNAs suppresses genes expressed in other cellular states (*Lee Lim-05 and Stephen M. Cohen-05*)
- Suppress cell-type specific genes

## 2) Control responses to changes in environment

- Protection against abnormal cellular states (p53 and NF- $\kappa$ B) and stabilize cellular states from transient “stress”

# Acknowledgments

Sharp Lab, MIT

Arvind Ravi

Joel Neilson

Grace Zheng

Anthony Leung

Margaret Ebert

Amanda Young

Mary Lindstrom

Allan Gurtan

Tyler Jacks

Madhu Kumar

Paul Chang

Sejal Vyas

Christopher Burge

Rickard Sandberg

Arup Sarma

Rudolf Jaenisch

Paul Anderson (HMS)

Nancy Kedersha

Rod Bronson (Tufts)



**KOCH INSTITUTE**  
for Integrative Cancer Research at MIT



Thank you for  
the opportunity  
to present this  
lecture